

ENVIRONMENT-DEPENDENT FUNDAMENTAL PHYSICAL CONSTANTS

–In the Theory of General Inconstancy *

Hidezumi Terazawa [†]

Center of Asia and Oceania for Science(CAOS)

3-11-26 Maesawa, Higashi-kurume, Tokyo 203-0032, Japan

and

Midlands Academy of Business & Technology(MABT),

Mansion House, 41 Guildhall Lane, Leicester LE1 5FR, United Kingdom

(Dedicated to the late Dr. Akira Tonomura who passed away on May 2, 2012)

Abstract

A theory of special inconstancy, in which some fundamental physical constants such as the fine-structure and gravitational constants may vary, is proposed in pregeometry. In the special theory of inconstancy, the $\alpha-G$ relation of $\alpha = 3\pi/[16 \ln(4\pi/5GM_W^2)]$ between the varying fine-structure and gravitational constants (where M_W is the charged weak boson mass) is derived from the hypothesis that both of these constants are related to the same fundamental length scale in nature. Furthermore, it leads to the prediction of $\dot{\alpha}/\alpha = (-0.8 \pm 2.5) \times 10^{-14} \text{yr}^{-1}$ from the most precise limit of $\dot{G}/G = (-0.6 \pm 2.0) \times 10^{-12} \text{yr}^{-1}$ by Thorsett, which is not only consistent with the recent observation of $\dot{\alpha}/\alpha = (0.5 \pm 0.5) \times 10^{-14} \text{yr}^{-1}$ by Webb *et al.* but also feasible for future experimental tests. In special inconstancy, the past and present of the Universe are explained and the future of it is predicted, which is quite different from that in the Einstein theory of gravitation. Also, a theory of general inconstancy, in which any fundamental physical constants may vary, is proposed in “more general relativity”, by assuming that the space-time is “environment-dependent”. In the general theory of inconstancy, the $G - \Lambda$ relation between the varying gravitational and cosmological

*A contribution to be presented at the Eighth Bolyai-Gauss-Lobachevsky International Conference on Non-Euclidean Geometry in Modern Physics and Mathematics, Uzhgorod, Ukraine, May 22- 25, 2012, to be published in the Proceedings, edited by L.L.Jenkovszky et al.(Bogolyubov Institute for Theoretical Physics, Kiev, 2012). This is the extended version of the previous invited talk presented at the Fifth International Conference “Methods of Non-Euclidean Geometry in Modern Physics”, Minsk, Belarus, October 10-13, 2006, in the Proceedings, edited by Yu.Kurochkin and V.Red’kov(Institute of Physics, National Academy of Sciences of Belarus, Minsk, 2006), p.20, which was the extended version of the recent lecture presented at the 12th International Seminar “Nonlinear Phenomena in Complex Systems”, Minsk, May 17-20, 2005, in the Proceedings, edited by L.Babichev and V.Kuvshinov (Institute of Physics, Minsk, 2005), Nonlin.Phenom.Complex Syst.**8**, 342(2005), which had been the extended version of the previous invited talk presented at the Fourth International Conference “Non-Euclidean Geometry in Modern Physics and Mathematics (BGL-4)”, Nizhny Novgorod, September 7-11, 2004, in the Proceedings, edited by L.L.Jenkovszky and G.M.Polotovskiy (Bogolyubov Institute for Theoretical Physics, Kiev, 2004), p.218.

[†]E-mail address: terazawa@mrj.biglobe.ne.jp

constants is derived from the hypothesis that the space-time metric is a function of τ , the “environment coordinate”, in addition to x^μ , the ordinary space-time coordinates. Furthermore, it leads to the prediction of the varying cosmological constant, which is consistent with the present observations. In addition, the latest observation of spatial variation in the fine-structure constant from VLT/UVES of $(1.1 \pm 0.2) \times 10^{-6} G\text{Ly}^{-1}$ by King *et al.* is suggested to be taken as a clear evidence for environment-dependent fundamental physical constants.

I. Introduction

Is a physical constant really constant? In 1937, Dirac[1] discussed possible time variation in the fundamental constants of nature. He made not only the large number hypothesis(LNH) but also, as a consequences of the LNH, the astonishing prediction that the gravitational constant G varies as a function of time. Since then, Jordan[2] and many others[3,4] have tried to construct new theories of gravitation or general relativity in order to accomodate such a time-varying G . Although the LNH has been inspiring many theoretical developements and has recently led myself[5] to many new large number relations, the prediction of the varying G has not yet received any experimental evidence. Recently, Thorsett[6] has shown that measurements of the masses of young and old neutron stars in pulsar binaries lead to the most precise limti of

$$\dot{G}/G = (-0.6 \pm 2.0) \times 10^{-12} yr^{-1}$$

at the 68% confidence level.

More recently, on the other hand, Webb *et al.*[7] have investigated possible time variation in the fine structure constant α by using quasar spectra over a wide rage of epoches, spanning redshifts $0.2 < z < 3.7$, in the history of our Universe, and derived the remarkable result of

$$\dot{\alpha}/\alpha = (6.40 \pm 1.35) \times 10^{-16} yr^{-1}$$

for $0.2 < z < 3.7$, which is consistent with a time-varying α . Note, however, that in 1976 Shylakhter[8] obtained the very restrictive limit of $|\Delta\alpha/\alpha| < 10^{-7}$ or, more precisely,

$$\Delta\alpha/\alpha\Delta t = (-0.2 \pm 0.8) \times 10^{-17} yr^{-1}$$

for $z \sim 0.16$ (but over a narrower and latest range of epochs between now and about 1.8 billion years ago) from the “Oklo natural reactor”. Very lately, Srianand *et al.*[9] have made a detailed many-multiplet analysis performed on a new sample of Mg II systems observed in high quality quasar spectra obtained using the Very Large Telescope and found a null result of $\Delta\alpha/\alpha = (-0.06 \pm 0.06) \times 10^{-5}$ for the fractional change in α or 3σ constraint of

$$-2.5 \times 10^{-16} yr^{-1} \leq (\Delta\alpha/\alpha\Delta t) \leq +1.3 \times 10^{-16} yr^{-1}$$

for $0.4 \leq z \leq 2.3$, which seems to be inconsistent with the result of Webb *et al.*[7]. However, a careful comparison of these different results[7-8] indicates that they are all consistent with a time-varying α as

$$\dot{\alpha}/\alpha = (0.5 \pm 0.5) \times 10^{-14} yr^{-1}$$

for $2.2 < z < 3.7$. More lately, Kanekar *et al.* have constrained the fundamental constant evolution with HI and OH lines as $\Delta\alpha/\alpha = (-1.7 \pm 1.4) \times 10^{-6}$ over a lookback time of 6.7Gyrs or, equivalently,

$$\dot{\alpha}/\alpha = (-2.5 \pm 2.1) \times 10^{-16} yr^{-1}$$

for $0 \leq z \leq 0.765$ [8].

In this talk, I am going to propose a theory of special inconstancy, in which some fundamental physical constants such as the fine-structure and gravitational constants may vary. In the special theory of inconstancy, the $\alpha - G$ relation of

$$\alpha = 3\pi/[16 \ln(4\pi/5GM_W^2)]$$

(where M_W is the charged weak boson mass) is derived from the hypothesis that both of α and G are related to the same fundamental length scale in nature. Furthermore, from the limit on $\dot{G}$ by Thorsett, it leads to the prediction of

$$\dot{\alpha}/\alpha = (-0.8 \pm 2.5) \times 10^{-14} yr^{-1}$$

which is not only consistent with the above result on $\dot{\alpha}$ but also feasible for future experimental tests. In special inconstancy, the past and present of the Universe are explained and the future of it is predicted, which is quite different from that in the Einstein theory of gravitation. Also, a general theory of inconstancy, in which any fundamental physical constants may vary, is proposed in “more general relativity”, by assuming that the space-time is “environment-dependent”. In the general theory of inconstancy, the “ $G - \Lambda$ relation” between the varying gravitational and cosmological constants is derived from the hypothesis that the space-time metric is a function of τ , the “environment coordinate”, in addition to x^μ , the ordinary space-time coordinates. Furthermore, it leads to the prediction of the varying cosmological constant, which is consistent with the present experimental observations.

I will organize this talk as follows: in Section II, I will briefly review pregeometry in which a theory of special inconstancy is constructed. In Sections III and IV, I will present the theories of special and general inconstancy and their predictions, respectively. Finally in Section V, I will discuss future prospects.

II. Pregeometry

Pregeometry is a theory in which Einstein geometrical theory of gravity in general relativity can be derived from a more fundamental principle as an effective and approximate theory at low energies (or at long distances). In 1967, Sakharov[10] suggested

possible approximate derivation of the Einstein-Hilbert action from quantum fluctuations of matter. A decade later, we[11] demonstrated that not only Einstein theory of gravity in general relativity but also the standard model of strong and electroweak interactions in quantum chromodynamics and in the unified gauge theory can be derived as an effective and approximate theory at low energies from the more fundamental unified composite model of all fundamental particles and forces[12].

Let us explain what pregeometry means more explicitly in a simple model of

$$S_0 = \int d^4x \sqrt{-g} L_0(g_{\mu\nu}(x), A_\mu(x), \phi_i(x))$$

where $g_{\mu\nu}$ is the space-time metric, $g = \det(g_{\mu\nu})$, A_μ is an Abelian gauge field, and ϕ_i ($i = 1 \sim n$) are n complex scalar fields of matter with the charge e . The fundamental Lagrangian L_0 consists of the gauge-invariant kinetic terms of the matter fields only as

$$L_0 = g^{\mu\nu} [\partial_\mu + iA_\mu] \phi_i^\dagger (\partial_\nu - iA_\nu) \phi_i - F^{-1}$$

(where F is an arbitrary constant) but does not contain either the kinetic term of the space-time metric or that of the gauge field so that both of $g_{\mu\nu}$ and A_μ are auxiliary fields. The effective action for the space-time metric and gauge field can be defined by the path-integral over the matter fields as

$$\exp(iS_{eff}) = \int \prod_i [d\phi_i^\dagger][d\phi_i] \exp(iS_0)$$

and it can be expressed formally as

$$S_{eff} = -i \text{Tr} \ln[(\partial_\nu - iA_\nu) \sqrt{-g} g^{\mu\nu} (\partial_\mu + iA_\mu)] - \int d^4x \sqrt{-g} F^{-1}$$

after the path-integration over ϕ_i . For small scalar curvature R and Ricci curvature tensor $R_{\mu\nu}$, the effective action can be calculated to be

$$S_{eff} = \int d^4x \sqrt{-g} [2\lambda + (1/16\pi G)R + c(R^2 + dR^{\mu\nu} R_{\mu\nu}) + (1/4e^2)F^{\mu\nu} F_{\mu\nu} + \dots]$$

with

$$2\lambda = [n\Lambda^4/8(4\pi)^2] - F^{-1},$$

$$(1/16\pi G) = n\Lambda^2/24(4\pi)^2,$$

$$c = n\ln\Lambda^2/240(4\pi)^2,$$

$$d = 2,$$

and

$$(1/4e^2) = n\ln\Lambda^2/3(4\pi)^2,$$

where λ and Λ are the cosmological constant and the momentum cut-off of the Pauli-Villars type, respectively. Note that the arbitrary constant F^{-1} plays a role of counter

term so that the cosmological constant may become as small as it is observed. Note also that the momentum cut-off Λ must be of order of the Planck mass $G^{-1/2}(\sim 10^{19}GeV)$. Furthermore, not only the R^2 and $R^{\mu\nu}R_{\mu\nu}$ terms but also the remaining terms in the expansion of S_{eff} are practically negligible. This complete a simple demonstration that not only the Einstein-Hilbert action of gravity but also the Maxwell action of electromagnetism in general relativity can be derived as an effective and approximate theory at low energies from the simple model in pregeometry, provided that there exists a natural momentum cut-off at around the Planck mass in nature[13].

One of the most remarkable consequences of pregeometry is the $\alpha - G$ relation, a simple relation between the fine-structure and gravitational constant, which can be easily derived from the results for α and for G by eliminating the momentum cut-off Λ . In our unified quark-lepton model of all fundamental forces[14,15], the $\alpha - G$ relation is given by[16]

$$\alpha = 3\pi / \sum_i Q_i^2 \ln(12\pi/nGm_i^2),$$

where Q_i and m_i are the charge and mass of quarks and leptons, respectively. For three generations of quarks and leptons and their mirror- or super-partners, the $\alpha - G$ relation simply becomes

$$\alpha \cong 3\pi/16 \ln(4\pi/5GM_W^2)$$

where M_W is the charged weak boson mass. Note that this $\alpha - G$ relation is very well satisfied by the experimental data of $\alpha \cong 1/137$, $G^{-1/2} \cong 1.22 \times 10^{19}GeV$, and $M_W \cong 80.4GeV$ [17].

III. Special Inconstancy

Special inconstancy is a principle in which some fundamental physical constants such as the fine-structure and gravitational constants may vary. Let us first make it clear that in this talk we use the natural unit system of $\hbar/2\pi = c = 1$ (where $\hbar$ is the Planck constant and c is the speed of light in vacuum). Note, however, that it does not mean that in discussing the relevant possibility of the varying fine-structure and gravitational constants[18], we exclude another intriguing possibility of the varying light velocity recently discussed by some authors[19] since varying either $\hbar$ or c is inevitably related to varying the fine-structure constant $\alpha (\equiv e^2/2\hbar c)$ (if the unit charge e stays constant). It simply means that we must set up a certain reference frame on which we can discuss whether physical quantities such as the fine-structure, gravitational, and cosmological[20] constants be really constant. Our basic hypothesis is that both of the fine-structure and gravitational constants are related to the more fundamental length scale of nature as in the unified (pregauge[21] and) pregeometric[10-12] theory (or “pregaugeometry” in short) of all fundamental forces[14,15] reviewed in the last Section.

To be more explicit, in the simple model of pregaugeometry discussed in the last Section, assert that

$$< (\partial_\mu + iA_\mu) \sqrt{-g} g^{\mu\nu} (\partial_\nu - iA_\nu) \phi_i >_\Lambda = 0,$$

$$g_{\mu\nu} = F < [(\partial_\mu + iA_\mu)\phi_i^\dagger](\partial_\nu - iA_\nu)\phi_i >_\Lambda,$$

and

$$A_\mu = (i/2) < [\phi_i^\dagger \partial_\mu \phi_i - (\partial_\mu \phi_i^\dagger) \phi_i] / (\phi_j^\dagger \phi_j) >_\Lambda,$$

where $<>_\Lambda$ denotes the expectation value in the space-time with the fundamental length scale parameter of Λ^{-1} . The first equation is the usual field equation for ϕ_i while the last two can be taken either as the “equations of motion” for $g_{\mu\nu}$ and A_μ , which can be derived from the fundamental action S_0 , or as the “fundamental field equations”, which can reproduce the effective Einstein-Hilbert-Maxwell action S_{eff} at low energies ($\ll \Lambda$) or at long distances ($\gg \Lambda^{-1}$).

The most important consequence of special inconstancy in pregaugeometry is the $\dot{\alpha} - \dot{G}$ relation for the varying fine-structure and gravitational constant of

$$\dot{\alpha}/\alpha^2 = (16/3\pi)[(\dot{G}/G) + 2(\dot{M}_W/M_W)],$$

which can be derived from differentiating both hand sides of the $\alpha - G$ relation with respect to any parameter for varying fundamental physical constants. This immediately leads to the remarkable predictions of

$$\dot{\alpha}/\alpha = (-0.8 \pm 2.5) \times 10^{-14} yr^{-1}$$

for constant M_W and

$$\dot{M}_W/M_W = (0.5 \pm 1.2) \times 10^{-12} yr^{-1}$$

from the limit of $\dot{G}/G = (-0.6 \pm 2.0) \times 10^{-12} yr^{-1}$ by Thorsett[6] and from the experimental data of $\dot{\alpha}/\alpha = (0.5 \pm 0.5) \times 10^{-14} yr^{-1}$ by Webb *et al.*[7]. The first prediction is not only consistent with the experimental data by Webb *et al.*[7] but also feasible for future experimental tests. The second prediction, however, seems too small to be feasible for experimental tests in the near future although such prediction for the possible varying particle masses seems extremely interesting at least theoretically. Note that the varying M_W is perfectly possible through the varying electroweak gauge coupling constant g (which is related to the fine-structure constant in the standard unified electroweak gauge theory of Glashow-Salam-Weinberg[22]) and/or the varying vacuum expectation value of the Higgs scalar v (which is related to the momentum cut-off Λ in the unified composite model of the Nambu-Jona-Lasinio type for all fundamental forces[23]) since $M_W = gv/2$. We must mention, however, that neither one of these predictions may assert for the fine-structure constant α or the weak boson mass M_W to vary since both of these predicted values are consistent with zero within their errors.

Let us now first add that in some pregaugeometric model[24], the $\alpha - G$ relation is not of the type of $\alpha \sim 1/\ln(1/GM^2)$ but of the type of $\alpha \sim GM^2$ (where M is a parameter of mass dimension) so that the $\dot{\alpha} - \dot{G}$ relation becomes

$$\dot{\alpha}/\alpha = (\dot{G}/G) + 2(\dot{M}/M).$$

This type of relation predicts

$$\dot{\alpha}/\alpha = (-0.6 \pm 2.0) \times 10^{-12} yr^{-1}$$

for constant M and

$$\dot{M}/M = (0.3 \pm 1.0) \times 10^{-12} yr^{-1}$$

from the limit by Thorsett[6] and from the experimental data by Webb *et al.*[7]. We suspect, however, that the predicted value for $\dot{M}/M$ seems too small to be feasible for experimental tests in the near future although the one for $\dot{\alpha}/\alpha$ is consistent with the experimental data by Webb *et al.*[7]. We must also mention that both of these predicted values are consistent with zero within their errors so that neither the fine-structure constant α nor the mass parameter M may be predicted to vary.

Next, remember that in the principle of special inconstancy we do not assert that physical constants may vary as a function of time but do that they may vary in general, depending on any parameters including the cosmological time, temperature, *etc.*. What is the origin of varying the physical constants? The answer to this question may be related to the answer to another fundamental question: What is the origin of the fundamental length scale Λ^{-1} in nature? It can be spontaneous breakdown of scale-invariance in the Universe, which has been proposed by myself[5] for the last quarter century. It can be the natural, dynamical, automatic, *a priori*, but somewhat “wishful-thinking” cut-off at around the Planck length $G^{1/2}$ where gravity would become as strong as electromagnetism, which was suggested by Landau[13] in 1955. It can also be due to the Kaluza-Klein extra dimension[25], which is supposed to be compactified at an extremely small length scale of the order of $G^{-1/2}$ or at a relatively large length scale of the order of $1/TeV$ recently emphasized by Arkani-Hamed *et al.*[26]. It seems, however, the most natural and likely that the origin of the fundamental length comes from the substructure of fundamental particles including quarks, leptons, gauge bosons, Higgs scalars, *etc.*[27,28]. In the unified composite model of all fundamental particles and forces[28], the fundamental energy scale Λ in pregeometery can be related to some even more fundamental parameters such as the masses of subquarks, the more fundamental constituents of quarks and leptons, and the energy scale in quantum subchromodynamics, the more fundamental dynamics confining subquarks into a quark or a lepton. In either way, the fundamental length scale Λ^{-1} can be identified with the size of quarks and leptons, the fundamental particles.

In pregeometric special inconstancy, let us now briefly explain the past and present of the Universe and try to predict the future of it, which may differ from that in the conventional Einstein theory of gravitation in general relativity[29]. The history of our Universe goes as follows: Long, long time ago there was no physical space-time, in which the space-time metric was finite and non-vanishing so that the distance was well defined, but the only matter “existed” in the mathematical space-time. Suddenly, there appeared the big bang of our Universe as a phase transition of the space-time from the pregeometric phase to the geometric one due to quantum fluctuations of matter, as suggested by us[30] in the early nineteen eighties, and our Universe had happened to be either flat or open. Then, not only all fundamental particles but also

all fundamental forces between them were created and they started obeying the effective theory of all fundamental particles and forces including the Einstein theory of gravity with the non-vanishing and varying cosmological constant. In the earliest era during which the matter density had been extremely small, our Universe had been expanding almost exponentially. It had been the “almost inflationary Universe”. In the next era of the radiation dominated Universe, our Universe was expanding less fast. Furthermore, in the last era of the matter dominated Universe, our Universe has still been expanding even faster. This history of our Universe is well simulated by a simple model of $(\Omega_m, \Omega_\lambda, -q) = (0, 1, 1)$, $(1/4, 3/4, 1/3)$, or $(1/4, 3/4, 1/2)$ for the early inflationary era, for the radiation dominated era, or for the matter dominated era, respectively, where Ω_m , Ω_λ , and q are the “pressureless-matter-density”, “scaled cosmological constant”, and deceleration parameter of the Universe, respectively. Note that there must be another “phase transition” in which Ω_λ changed from 1 to $2/3$ in between the early inflationary era and the radiation dominated era. Concerning the cosmological constant, I have been most impressed by the recent observation of the “farthest supernova ever seen” by Hubble Space Telescope[31]. “This supernova shows us the universe is behaving like a driver who slows down approaching a red stoplight and then hits the accelerator when the light turns green.” Note that this behavior of the Universe is what our model simulates. Note also that our model of the Universe is consistent with the recent measurement of the cosmological mass density from clustering in the Two-Degree-Field Galaxy Redshift Survey[32] which strongly favors a low density Universe with $\Omega_m \cong 0.3$. Very recently, the CBI Collaboration[33,17] has found $\Omega_\lambda = 0.64 + 0.11 / - 0.14$ and $\Omega_m + \Omega_\lambda = 1.006 \pm 0.006$. More recently, the Wilkinson Microwave Anisotropy Probe (WMAP) team[34,17] has found 1) the first generation of stars to shine in the Universe first ignited only 200 million years after the big bang, 2) the age of the Universe is 13.69 ± 0.13 billion years old, and 3) $\Omega_m = 0.26 \pm 0.02$ and $\Omega_\lambda = 0.74 \pm 0.03$.

The future of the Universe in our special inconstant picture can be quite different from that in the Einstein-Friedmann picture: 1) Since the cosmological constant may vary in special inconstancy, the space-time of our Universe which is almost flat and expanding faster and faster may not continue to be flat and accelerating forever. Our Universe may even encounter a “topological phase transition”, which was first discussed by Wheeler[35] in 1959, from the open Universe to the closed one. 2) If the gravitational constant increases, the expansion of the space-time may not continue forever. The Universe may well stop expanding, start contracting, and even be bouncing forever. If G decreases, it will be more accelerated ever. 3) If the fine-structure constant (and/or other fundamental coupling constants such as the strong and weak coupling constants) varies, our Universe may encounter as “obsolete phase transition” from the matter-dominated Universe to the radiation-dominated one. In short, we can expect anything about the future of our Universe or, in other words, we can predict nothing definite on the destiny of our Universe.

IV. General Inconstancy

General inconstancy is a principle in which any fundamental physical constants may vary. In order to accomodate general inconstancy with general relativity, let us assume that the space-time is environment dependent. More explicitly, let us make the hypothesis that the space-time metric, $g_{\mu\nu}$, is a function of the “environment coordinate”, τ , in addition to the ordinary space-time coordinates, x^μ , where $\mu, \nu = 0, 1, 2, 3$, so that the infinitesimal distance, ds , is given by

$$ds^2 = g_{\mu\nu}(x, \tau) dx^\mu dx^\nu.$$

Note that the physical meaning of τ is arbitrary as it can be the temperature(T), the cosmological time(t), a parameter related to the Kaluza-Klein extra dimensions, or anything else. Note also that τ may not be a continuous valuable but be a discrete number and that there may exist more than one “environment coordinates”. I will consider such extensions of the environment-dependent space-time metric later.

In this “more general relativity”, let us consider a simple model for general inconstancy given by the generalized Einstein-Hilbert action of

$$S_I = \int d\tau \int d^4x \sqrt{-g} [R(x, \tau) + \dots] = \int d^4x \sqrt{-g} [2\lambda + (1/16\pi G)R(x) + \dots].$$

In this equation we may find the equations of

$$2\lambda = \int d\tau \int d^4x \sqrt{-g} [R(x, \tau) + \dots] / \int d^4x \sqrt{-g}$$

and

$$(1/16\pi G) = \int d\tau \int d^4x \sqrt{-g} [R(x, \tau) + \dots] / \int d^4x \sqrt{-g} R(x)$$

so that we may obtain the $G - \Lambda$ relation of

$$32\pi G\lambda = \int d^4x \sqrt{-g} R(x) / \int d^4x \sqrt{-g}.$$

Futhermore, by differentiating both hand-sides of the relation with respect to τ , we can obtain the relation between the varying gravitaional and cosmological constants:

$$\dot{G}/G = -\dot{\lambda}/\lambda.$$

Since in the homogeneous and isotropic Einstein-Friedmann model of the Universe, the scalar curvature is given by $R = 6[(1-q)H^2 + (k/a^2)]$, where a is the scale parameter in the Robertson-Walker metric and $k = +1, 0, -1$ for the closed, flat, or open universe, the first relation explains the positivity of the cosmological constant, λ , for the present Universe with $-q > 0$ and, probably, $k = 0$. The second relation leads to the prediction of

$$\dot{\lambda}/\lambda = (0.6 \pm 2.0) \times 10^{-12} yr^{-1},$$

from the limit by Thorsett[6]. This prediction is not only consistent with the present experimental observation but also feasible for future experimental tests.

I must mention, however, that I have not completed checking whether all the possible consequences of this simple theory of general inconstancy be consistent with all the presently available experimental observations. It is certainly the most important subject for future investigations. I have not found either whether it is necessary to extend it for discrete τ or for more than one “environment coordinates”. Just imagine that we stand in front of the entrance gate for the vast unknown world of general inconstancy!

V. Future Prospects

In conclusion, let us point out that not only continuous physical constants such as α and G but also discrete physical numbers such as the number of the space-time dimensions n , the number of quark colors N_c , the number of quark-lepton generations N_g , *etc.* may vary. In fact, an astonishing “dimensional phase transition”, which was discussed by myself[36] about two decades ago, may be possible in the history of our Universe. If n is related to N_c as in the “space-color correspondence”, which was proposed by myself about three decades ago[37], both of these fundamental physical natural numbers must vary simultaneously. Before concluding this talk, let me ask the following question: Are no constants of nature constant? After all, it may be that nothing is constant or permanent in the Universe as emphasized by the Greek and Indian philosophers about two and a half millennia ago!

Recently, the OPERA neutrino experiment[38] at the underground Gran Sasso Laboratory has measured the velocity of neutrinos from the CERN CNGS beam over a baseline of about 730 km and claimed that a relative difference of the muon neutrino velocity with respect of the speed of light $(v - c)/c = (2.48 \pm 0.24(stat.) \pm 0.30(sys.)) \times 10^{-5}$. This once intriguing observation of the superluminal muon neutrinos, if it were real, might be taken as an evidence for possible violation of special relativity due to the environment-dependent velocity of light in vacuum in theory of general-inconstancy.

In ending this lecture, let us introduce two more recent reports on finding an evidence for environment-dependent fundamental physical constants in addition to many recent reports[39]. Very lately, Kanekar *et al.* have obtained a very strong constraint of $\Delta\alpha/\alpha = (-1.7 \pm 1.4) \times 10^{-6}$ over a lookback time of 6.7Gyrs with HI and OH lines[40]. Even more lately, King *et al.* have found spatial variation of the fine-structure constant with the amplitude of $(1.1 \pm 0.2) \times 10^{-6} GLyr^{-1}$ by using UVES on the VLT[41], which may be taken as a clear evidence for the environment-dependent fundamental physical constant. It may be a big-bang in physics!

Acknowledgement

The author would like to thank Professor Laszlo L. Jenkovszky, and all the other organizers of this Eighth International Conference on Methods of Non-Euclidean Geometry in Modern Physics, Uzhgorod, Ukraine, May 22- 25, 2012 for inviting him to the Conference, although he could not attend the Conference, and for publishing this contribution in the Proceedings. This contribution has been dedicated to Dr. Akira Tonomura, my late friend, who first observed the Aharonov-Bohm effect.

References

- [1] P.M.A.Dirac, Nature **139**, 323(1937); Proc.R.Soc.London **A165**, 199(1938); **A333**, 403(1973); **A338**, 439(1974).
- [2] P.Jordan, Ann.der Phys. **6**, Folge **1**, 219(1947). For the recent related article, see E.L.Schuking, Physic today, October 1999, vol.**52**, No. 10, p.26(1999).
- [3] F.Hoyle and J.V.Narlikar, Nature **233**, 41(1971).
- [4] V.Canuto, P.J.Adams, S.-H.Hsieh, and E.Tsiang, Phys.Rev.D **16**, 1643(1977).
- [5] H.Terazawa, Phys.Lett. **101B**, 43(1981); in **Proc. 3rd Alexander Friedmann International Seminar on Gravitation and Cosmology**, St.Petersburg, 1995, edited by Yu.N.Gnedin, A.A.Grib, and V.M.Mostepanenko (Friedmann Laboratory Pub., St.Petersburg, 1999), p.116; Mod.Phys.lett. **A11**, 2971(1996); **A12**, 2927(1997); **A13**, 2801(1998).
- [6] S.E.Thorsett, Phys.Rev.Lett. **77**, 1432(1996). See also V.M.Kaspi, J.H.Taylor, and M.F.Ryba, Astrophys.J. **42**, 713(1996) and D.B.Guenther, L.M.Krauss, and P.Demarque, *ibid.***498**, 871(1998). For a recent review, see J.Uzan, Rev.Mod.Phys. **75**, 403(2003). More recently, Copi *et al.* have inferred $-4 \times 10^{-13} yr^{-1} < (\dot{G}/G)_{today} < 3 \times 10^{-13} yr^{-1}$ (by assuming a monotonic power law time dependence $G \propto t^{-a}$) from the big-bang nucleosynthesis constraint. See C.J.Copi, A.N.Davis, and L.M.Krauss, Phys.Rev.Lett. **92**, 171301(2004). Very lately, Jofré *et al.* have estimated one of the most restrictive upper limits of $|\dot{G}/G| < 4 \times 10^{-12} yr^{-1}$ by using the surface temperature of the nearest millisecond pulsar, PSR J0437-4715, (inferred from ultraviolet observations) and by considering only modified Urca reactions. See P.Jofré, A.Reisenegger, and R.Fernández, Phys.Rev.Lett. **97**, 131102(2006).
- [7] J.K.Webb *et al.*, Phys.Rev.Lett. **82**, 884(1999); *ibid.***87**, 091301(2001); M.T.Murphy, J.K.Webb, and V.V.Flambaum, Mon.Not.R.Astron.Soc. **345**, 609(2003). Recently, de Souza has claimed that the velocity in which quasars evolve into normal galaxies make us to see the limit of quasars slightly Doppler shifted which explains the “misleading” variation of the fine structure constant. See M.E.De Souza, astro-ph/0301085, 6 Jan 2003. However, it seems that his estimate of the shift is overestimated by ignoring the isotropy of quasar expansions. Also recently, J.N.Bahcall *et al.* have set a robust upper limit on $\dot{\alpha}$, $\dot{\alpha}/\alpha < 10^{-13} yr^{-1}$, by using the strong nebular emission lines of O III. See J.N.Bahcall, C.L.Steinhardt, and D.Schlegel, Astrophys.J. **600**, 520(2004). See also L.L.Cowie and A.Songaila, Astrophys.J. **453**, 596(1995); R.L.White, R.H.Becker, X.Fan, and M.A.Strauss, Astron.J. **126**, 1(2003); A.Songaila, astro-ph/0402347, to be published in Astron.J.. For a recent review, see A.Songaila and L.L.Cowie, Nature **398**, 667(1999). Very recently, Tzanavaris *et al.* have estimated the time variation of $x(= \alpha^2 g_p \mu$ where g_p is the proton g factor and μ is the electron/proton mass ratio, m_e/m_p) from 21-cm

and ultraviolet quasar absorption lines over a redshift range $0.24 < z < 2.04$ as $\dot{x}/x = (-1.43 \pm 1.27) \times 10^{-15} \text{yr}^{-1}$ for a linear fit. Using the $\Delta\alpha/\alpha$ results and assuming $\Delta g_p/g_p \sim 0$, they have obtained $\Delta\mu/\mu = (2.31 \pm 1.03) \times 10^{-5}$ and $\Delta\mu/\mu = (1.29 \pm 1.01) \times 10^{-5}$. See P.Tzanavaris *et al.*, Phys.Rev.Lett. **95**, 041301(2005). More recently, Reinhold *et al.* have obtained $\Delta\mu/\mu = (2.4 \pm 0.6) \times 10^{-5}$ for a weighted fit and $\Delta\mu/\mu = (2.0 \pm 0.6) \times 10^{-5}$ for an unweighted fit, which indicates, at a 3.5σ confidence level, that μ could have decreased in the past 12 Gyr. See E.Reinhold *et al.*, Phys.Rev.Lett. **96**, 151101(2006). Note, however, that the possible variation of μ may not necessarily mean that of the fundamental mass scale in nature but simply do that of the effective electron and/or proton masses due to different environmental parameters such as density, temperature, *etc.*.

- [8] A.I.Shylakhter, Nature **264**, 340(1976); ATOMKI Report No.A/1(1983); T.Damour and F.Dyson, Nucl.Phys. **B480**, 37(1996); Y.Fujii *et al.*, *ibid.***B573**, 377(2000), hep-ph/0205206. For the latest reexamination of the Oklo and other constrains, see Y.Fujii and A.Iwamoto, Phys.Rev.Lett. **91**, 241302(2003). Very recently, Marion *et al.* have set a stringent upper bound on $\dot{\alpha}$, $\dot{\alpha}/\alpha = (-0.4 \pm 16) \times 10^{-16} \text{yr}^{-1}$, by comparing the hyperfine frequencies of Cs and Rb atoms. See H.Marion *et al.*, Phys.Rev.Lett. **90**, 150801(2003). See also S.Bize *et al.*, *ibid.***90**, 150802(2003); J.N.Chengalur and N.Kanekar, *ibid.***91**, 241302(2003); T.Ashenfelter, G.J.Mathews, and K.A.Olive, *ibid.***92**, 041102(2004). Even more recently, Fisher *et al.* have deduced a seperate limit of $\dot{\alpha}/\alpha = (-0.9 \pm 2.9) \times 10^{-15} \text{yr}^{-1}$ from combining the $1S - 2S$ transition frequency in H atom with the optical transition frequency in Hg^+ against the hyperfine splitting in Cs and the microwave Rb and Cs clock comparison. See M.Fisher, Phys.Rev.Lett. **92**, 230802(2004). Very lately, Peik *et al.* have measured an optical transition frequency at 688THz in $^{171}\text{Yb}^+$ with a cesium atomic clock at 2 times separated by 28yr and find a value for the fractional variation of the freauency ratio $f_{\text{Yb}}/f_{\text{Cs}}$ of $(-1.2 \pm 4.4) \times 10^{-15} \text{yr}^{-1}$, consistent with zero. Combined with recently published values for the constancy of other transition frequencies this measurement sets an upper limit on the present variability of α at the level of $2.0 \times 10^{-15} \text{yr}^{-1} (1\sigma)$ corresponding so far to the most stringent limit from laboratory experiments. See E.Peik *et al.*, Phys.Rev.Lett. **93**, 170801(2004). For a recent brief review, see C.Seife, Science **306**, 793(2004).
- [9] H.Chand, R.Srianand, P.Petitjean, and B.Arakil, Astophs. **417**, 853(2004); R.Srianand *ibid.*, Phys.rev.Lett. **92**, 121302(2004). For a latest comparison of this data with the Oklo data and the quasar data, see L.L.Cowie and A.Songaila, Nature **428**, 132(2004).
- [10] A.D.Sakharov, Dokl.Akad.Nak SSSR **177**, 70(1967)[Sov.Phys.JETP **12**, 1040(1968)].
- [11] K.Akama, Y.Chikashige, T.Matsuki, and H.Terazawa, Prog.Theor.Phys. **60**, 868(1978); S.L.Adler, Phys.Rev.Lett. **44**, 1567(1980); A.Zee, Phys.Rev.D **23**, 858(1981); D.Amati and G.Veneziano, Phys.Lett. **105B**, 358(1981).

- [12] For a review, see, for example, H.Terazawa, in **Proc. 1st A.D.Sakharov Conference on Physics**, Moscow, 1991, edited by L.V.Keldysh and V.Ya.Fainberg (Nova Science, New York, 1992), p.1013.
- [13] L.Landau, in **Niels Bohr and the Development of Physics**, edited by W.Pauli(Mc Graaw-Hill, New York, 1955), p.52. In 1977, we suggested abnormally strong gravity at short distances which breaks some presently reliable basic principles in quantum field theories such as special relativity and microscopic causality at a distance of the order of the Planck length. We also suggested that baryon number generation occurred in the very early Universe, breaking CPT invariance which is based on these basic principles. See H.Terazawa, M.Kasuya, and K.Akama, Report No. INS-Report-331 (INS, Univ. of Tokyo, 1979); H.Terazawa, in **Proc. Workshop on Unified Theories and Baryon Number in the Universe**, Tsukuba, 1979, edited by O.Sawada and A.Sugamoto (National Lab. for High Energy Physics, Tsukuba, 1979), p.99; *Gen.Relativ.Gravit.* **12**, 93(1980). For a recent idea similar to the latter suggestion, *i.e.*, dynamical breaking of CPT in an expanding Universe and, combined with baryon-number-violating interactions, driving the Universe towards an equilibrium baryon asymmetry that is observationally acceptable, see H.Davoudiasl *et al.*, *Phys.Rev.Lett.* **93**, 201301(2004). For a very recent search for signatures of Lorentz and CPT violations, see B.Feng *et al.*, *Phys.Rev.Lett.* **96**, 221302(2006). Also, recently Calmet *et al.* have derived fundamental limits on measurements of position arising from quantum mechanics and classical general relativity and shown that any primitive probe or target used in an experiment must be larger than the Planck length. See X.Calmet, M.Graesser, and S.D.H.Hsu, *Phys.Rev.Lett.* **93**, 211101(2004). Very lately, Robinson and Wilczek have calculated the contribution of graviton exchange to the running of gauge couplings at lowest nontrivial order in perturbation theory and found that, when extrapolated formally, it renders all gauge couplings asymptotically free. See P.Robinson and F.Wilczek, *Phys.Rev.Lett.* **96**, 231601(2006). Their results may lead to a modified version of the Landau conjecture that the effect of gravity suppresses all gauge couplings at extremely short distances.
- [14] H.Terazawa, Y.Chikashige, and K.Akama, *Phys.Rev.D* **15**, 480(1977).
- [15] For reviews, see, for example, H.Terazawa, in **Proc. 19th International Conf. on High Energy Physics**, Tokyo, 1978, edited by S.Homma, M.Kawaguchi, and H.Miyazawa (*Phys.Soc.Japan*, Tokyo, 1979), p.617; in **Proc. 22nd International conf. on High energy Physics**, Leipzig, 1984, edited by A.Meyer and E.Wieczok (Akademie der Wissenschaften der DDR, Zeuthen, 1984), vol.I, p.63.
- [16] H.Terazawa, Y.Chikashige, K.Akama, and T.Matsuki, *Phys.Rev.D* **15**, 1181(1977); H.Terazawa, *ibid.* **16**, 2373(1977); **22**, 1037(1980); **41**, 3541(E)(1990).
- [17] K.Nakamura *et al.*(Particle Data Group), *J.Phys.G* **37**, 075021(2010).
- [18] J.D.Bekenstein, *Phys.Rev.D* **25**, 1528(1982); J.W.Moffat, *Int.J.Mod.Phys.* **D2**, 351(1992); J.D.barrow and J.Magueijo, *Phys.Lett.* **B443**, 104(1998).

- [19] M.A.Clayton and J.W.Moffat, Phys.Lett. **B460**, 263(1999); A.Albrecht and J.Magueijo, Phys.Rev.D **59**, 043516(1999); J.D.Barrow, *ibid.***59**, 043515(1999); P.P.Avelino and C.J.A.P.Martins, Phys.Lett. **B459**, 468(1999); P.C.W.Davies *et al.*, Nature **418**, 603(2002).
- [20] A.Vilenkin, Phys.Rev.Lett. **81**, 5501(1998); A.A.Starobinsky, JETP Lett. **68**, 757(1998); L.O.Pimental and L.M.Diaz-rivera, Int.J.Mod.Phys. **A14**, 1523(1999).
- [21] J.D.Bjorken, Ann.Phys. (N.Y.) **24**, 174(1963); H.Terazawa, Y.Chikashige, and K.Akama, Ref.[14].
- [22] S.L.Glashow, Nucl.Phys. **22**, 579(1961); A.Salam in **Elementary Particle Physics**, edited by N.Svartholm (Almqvist and Wilsell, Stockholm, 1968), p.367; S.Weinberg, Phys.rev.Lett. **19**, 1264(1967).
- [23] Y.Nambu and G.Jona-Lasinio, Phys.Rev. **122**, 345(1961); H.Terazawa, Y.Chikashige, and K.Akama, in Ref.[14].
- [24] See, for example, H.Terazawa, Phys.Lett. **133B**, 57(1983).
- [25] T.Kaluza, Sitzker.Preuss.Akad.Wiss. **K1**, 966(1921); O.Klein, Z.Phys. **37**, 896(1926).
- [26] N.Arkani-Hamed, S.Dimopoulos, and G.Dvali, Phys.Lett. **B429**, 263(1998); Phys.Rev.D **59**, 086004(1999); I.Antoniadis *et al.*, Phys.Lett. **B436**, 257(1998); L.Randall and R.Sundrum, Phys.Rev.Lett. **83**, 3370(1999); **83**, 4690(1999).
- [27] J.C.Pati and A.Salam, Phys.Rev.D **10**, 275(1974); H.Terazawa, Y.Chikashige, and K.Akama, in Ref.[14]; H.Terazawa, Phys.Rev.D **22**, 185(1980).
- [28] For classical reviews, see, for example, H.Terazawa, in Ref.[15] and for a latest review, see for example, H.Terazawa, in **Proc. International Conf. “New Trends in High-Energy Physics”**, Alushta, Crimea, 2003, edited by P.N.Bogolyubov, L.L.Jenkowszky, and V.V.Magas (bogolyubov Institute for Theoretical Physics, Kiev, 2003), Ukranian J.Phys. **48**, 1292(2003).
- [29] For the latest more detailed review including many other current topics in cosmology, see, for example, H.Terazawa, in **Proc. 11th Interational Conf. and School “Foundations & Advances in Nonlinear Science”**, Minsk, Belarus, 2003, edited by V.I.Kuvshinov and G.G.Krylov(Belarusian State University, Minsk, Belarus, 2004), p.99.
- [30] K.Akama and H.Terazawa, Gen.Rel.Grav. **15**, 201(1983).
- [31] For a latest review, see C.Seife, Science **303**, 1271(2004). Even more lately, an extremely small galaxy has been found with the image taken by the Very Large Telescope. For a brief review, see C.Seife, *ibid.***303**, 1597(2004).

- [32] J.A.Peacock *et al.*, Nature **410**, 169(2001). For a recent brief review, see, for example, A.Watson, Science **295**, 2341(2002).
- [33] J.L.Sievers *et al.* (CBI Collaboration), Astrophys.J. **591**, 599(2003).
- [34] C.L.Bennett *et al.*, Astrophys.J.Suppl. **148**, 1(2003); D.N.Spergel *et al.*, *ibid.* **148**, 175(2003). See also T.Reichhardt, Nature **421**, 777(2003); S.Carroll, *ibid.* **422**, 26(2003); G.Brumfiel *ibid.* **422**, 108(2003). for a review, see B.Schwarzschild, Physics Today, Vol.56, No.4, April 2003, p.21.
- [35] J.A.Wheeler, in **Conference on the Role of Gravitation in Physics**, Chapel Hill, North Carolina, 1959(WADC Technical report 57-216, ASTIA Document No. AD118180; B.DeWitt, in **Proc. Third Seminar on Quantum Gravity**, Moscow, 1984, edited by M.A.Markov, V.A.Berezin, and V.P.Frolov (World Scientific, Singapore, 1985), p.103; V.G.Gurzadyan and A.A.Kocharyan, Zh.Eksp.Teor.Fiz. **95**, 3(1989)[Sov.Phys.JETP **8**, 1(1989); Mod.Phys.Lett. **A4**, 50(1989).
- [36] H.Terazawa, in **Quantum field Theory and Quantum Statistics**, edited by I.A.Batalin and G.A.Vilkovisky (Adam Hilger, London, 1987), Vol.I, p.637; in **Particles and Nuclei**, edited by H.Terazawa (World Scientific, Singapore, 1986), p.304; in **Modern Problems of the Unified Field Theory**, edited by D.V.Golt'sov, L.S.Kuzmenkov, and P.I.Pronin (Moscow State University, Moscow, 1991), p.124 (in russian translation); in **Proc. 5th Seminar on Quantum Gravity**, Moscow, 1990, edited by M.A.Markov, V.A.Berezin, and V.P.Frolov (World Scientific, Singapore, 1991), p.643. For a recent similar idea of emergence of a four dimensional world from causal quantum gravity, see J.Ambjørn, J.Jurkiewicz, and R.Loll, Phys.Rev.Lett. **93**, 131301(2004).
- [37] H.Terazawa, in ref.[26]; in **Wandering in the Fields**, edited by K.Kawarabayashi and A.Ukawa (World Scientific, Singapore, 1987), p.388.
- [38] T.Adam *et al.*(OPERA Collaboration), arXiv:1109.4897v1[hep-ex], Thu, 22 Sep 2011.
- [39] For a recent review, see J.C.Berengut and V.V.Flambaum, Nuclear Physics News, Vol.20, No.3, p.19(2010).
- [40] N.Kanekar *et al.*, arXiv:1201.3372v1[astro-ph.CO], Mon, 16 Jan 2012, to be published in Astrophys.J.Letters(2012).
- [41] J.A.King *et al.*, arXiv:1202.4758v1[astro-ph.CO], Tue, 21 Feb 2012, to be published in Mon.Not.R.Astron.Soc.**000**, 1-38(2012).